

# ZERO MEAN CURVATURE SURFACES IN LORENTZ-MINKOWSKI 3-SPACE WHICH CHANGE TYPE ACROSS A LIGHT-LIKE LINE

S. FUJIMORI, Y. W. KIM, S.-E. KOH, W. ROSSMAN, H. SHIN, M. UMEHARA,  
K. YAMADA AND S.-D. YANG

ABSTRACT. It is well-known that space-like maximal surfaces and time-like minimal surfaces in Lorentz-Minkowski 3-space  $\mathbf{R}_1^3$  have singularities in general. They are both characterized as zero mean curvature surfaces. We are interested in the case where the singular set consists of a light-like line, since this case has not been analyzed before. As a continuation of a previous work by the authors, we give the first example of a family of such surfaces which change type across the light-like line. As a corollary, we also obtain a family of zero mean curvature hypersurfaces in  $\mathbf{R}_1^{n+1}$  that change type across an  $(n-1)$ -dimensional light-like plane.

## 1. INTRODUCTION

Many examples of space-like maximal surfaces containing singular curves in the Lorentz-Minkowski 3-space  $(\mathbf{R}_1^3; t, x, y)$  of signature  $(-++)$  have been constructed in [12], [1], [13], [9] and [4].

In this paper, we are interested in the zero mean curvature surfaces in  $\mathbf{R}_1^3$  changing their causal type: Klyachin [11] showed under a sufficiently weak regularity assumption that a zero mean curvature surface in  $\mathbf{R}_1^3$  changes its causal type only on the following two subsets:

- null curves (i.e., regular curves whose velocity vector fields are light-like) which are non-degenerate (cf. Definition 2.1), or
- light-like lines, which are degenerate everywhere.


Given a non-degenerate null curve  $\gamma$  in  $\mathbf{R}_1^3$ , there exists a zero mean curvature surface which changes its causal type across this curve from a space-like maximal surface to a time-like minimal surface (cf. [6], [11], [10] and [8]). This construction can be accomplished using the Björling formula for the Weierstrass-type representation formula of maximal surfaces. By unifying the results of Gu [6], Klyachin [11], and [8], we explain the mechanism for how zero mean curvature surfaces change type across non-degenerate null curves, and give ‘the fundamental theorem of type change for zero mean curvature surfaces’ (cf. Theorem 2.17) in the second section of this paper.

---

*Date:* November 21, 2012.

*Key words and phrases.* maximal surface, minimal surface, type change, zero mean curvature.

Kim was supported by NRF 2009-0086794, Koh by NRF 2009-0086794 and NRF 2011-0001565, and Yang by NRF 2012-042530. Fujimori was partially supported by the Grant-in-Aid for Young Scientists (B) No. 21740052, Rossman was supported by Grant-in-Aid for Scientific Research (B) No. 20340012, Umehara by (A) No. 22244006 and Yamada by (B) No. 21340016 from Japan Society for the Promotion of Science.

FIGURE 1. Hyperbolic catenoids  $\mathcal{C}_+$ ,  $\mathcal{C}_-$ .

However, if  $\gamma$  is a light-like line, the aforementioned construction fails, since the isothermal coordinates break down at the light-like singular points (cf. the proof of Theorem 2.17). Locally, such a surface is the graph of a function  $t = f(x, y)$  satisfying

$$(1.1) \quad (1 - f_y^2)f_{xx} + 2f_x f_y f_{xy} + (1 - f_x^2)f_{yy} = 0.$$

We call this and its graph the *zero mean curvature equation* and a *zero mean curvature surface*, respectively. Zero mean curvature surfaces containing light-like lines have not yet been sufficiently studied. In a previous work with Hidenobu Takahashi [3], the authors give several fundamental examples of such surfaces.

For example, the space-like hyperbolic catenoid

$$(1.2) \quad \mathcal{C}_+ = \{(t, x, y) \in \mathbf{R}_1^3; \sin^2 x + y^2 - t^2 = 0\}$$

and the time-like hyperbolic catenoid

$$(1.3) \quad \mathcal{C}_- = \{(t, x, y) \in \mathbf{R}_1^3; \sinh^2 x + y^2 - t^2 = 0\}$$

are both typical examples of zero mean curvature surfaces containing singular light-like lines as subsets (cf. Figure 1). The space-like hyperbolic catenoid  $\mathcal{C}_+$  is singly periodic.

Also, both the space-like Scherk surface (cf. [3])

$$(1.4) \quad \mathcal{S}_+ = \{(t, x, y) \in \mathbf{R}_1^3; \cos t = \cos x \cos y\}$$

and the time-like Scherk surface of first kind (cf. [3])

$$(1.5) \quad \mathcal{S}_- = \{(t, x, y) \in \mathbf{R}_1^3; \cosh t = \cosh x \cosh y\}$$

contain singular light-like lines as subsets (cf. Figure 2).  $\mathcal{S}_+$  is triply periodic.

As an application of the results in Section 2, we show in Section 3 that  $\mathcal{C}_+$  and  $\mathcal{C}_-$  induce a common zero mean curvature graph (cf. Figure 3, left)



$$(1.6) \quad \mathcal{C}_0 = \{(t, x, y) \in \mathbf{R}_1^3; t = y \tanh x\}$$

via their conjugate surfaces. Similarly, we also show that the Scherk-type surfaces  $\mathcal{S}_+$  and  $\mathcal{S}_-$  induce a common zero mean curvature graph via their conjugate surfaces (cf. Figure 3, right)

$$(1.7) \quad \mathcal{S}_0 = \{(t, x, y) \in \mathbf{R}_1^3; e^t \cosh x = \cosh y\}.$$

These two phenomena were briefly commented upon in [3].

On the other hand, zero mean curvature surfaces which actually change type across a light-like line were unknown. For example, the above four surfaces  $\mathcal{C}_\pm$  and

FIGURE 2. Scherk-type surfaces  $\mathcal{S}_+$  and  $\mathcal{S}_-$ .FIGURE 3. Zero mean curvature graphs  $\mathcal{C}_0$  and  $\mathcal{S}_0$  (the curves where the surfaces change type are also indicated).

$\mathcal{S}_\pm$  do not change type. As announced in [3], the main purpose of this paper is to construct such an example. In Section 4, we give a formal power series solution of the zero mean curvature equation describing all zero mean curvature surfaces which contains a light-like line. Using this, we give the precise statement of our main result and show how the statement can be reduced to a proposition (cf. Proposition 4.3). In Section 5, we then prove it. As a consequence, we obtain the first example of (a family of) zero mean curvature surfaces which change type across a light-like line.

At the end of this paper, we provide an appendix which prepares several inequalities used in Section 4.

## 2. TYPE CHANGE OF ZERO MEAN CURVATURE SURFACES

In this section, we discuss type changes for zero mean curvature surfaces by unifying the results of Gu [6], Klyachin [11] and four of the authors here [8].

A regular curve  $\gamma : (a, b) \rightarrow \mathbf{R}_1^3$  is called *null* or *isotropic* if  $\gamma'(t) := d\gamma(t)/dt$  is a light-like vector for all  $t \in (a, b)$ .

**Definition 2.1.** A null curve  $\gamma : (a, b) \rightarrow \mathbf{R}_1^3$  is called *degenerate* or *non-degenerate* at  $t = c$  if  $\gamma''(c)$  is or is not proportional to the velocity vector  $\gamma'(c)$ , respectively. If  $\gamma$  is non-degenerate at each  $t \in (a, b)$ , it is called a *non-degenerate* null curve.

We now give a characterization of zero mean curvature surfaces that change type across a real analytic non-degenerate null curve. Given an arbitrary real analytic null curve  $\gamma : (a, b) \rightarrow \mathbf{R}_1^3$ , we denote the unique analytic extension of it still by  $\gamma$  throughout this article, by abusing notation slightly. We consider the two surfaces

$$\Phi(u, v) := \frac{\gamma(u + iv) + \gamma(u - iv)}{2},$$

and

$$\Psi(u, v) := \frac{\gamma(u + v) + \gamma(u - v)}{2},$$

which are defined for  $v$  sufficiently close to zero. We recall the following assertion:

**Proposition 2.2** ([6], [11] and [8]). *Given a real analytic non-degenerate null curve  $\gamma : (a, b) \rightarrow \mathbf{R}_1^3$ , the union of the images of  $\Phi$  and  $\Psi$  given as above are subsets of a real analytic immersion, and the intersection is  $\gamma$ . Moreover,  $\Phi$  gives a space-like maximal surface and  $\Psi$  gives a time-like minimal surface if  $v$  is sufficiently close to zero. Furthermore, this analytic extension of the curve  $\gamma$  as a zero mean curvature surface does not depend upon a choice of the real analytic parametrization of the curve  $\gamma$ .*

*Proof.* We give here a proof for the sake of readers' convenience. We have that

$$\Phi(u, v) = \sum_{n=0}^{\infty} (-1)^n \gamma^{(2n)}(u) v^{2n}, \quad \Psi(u, v) = \sum_{n=0}^{\infty} \gamma^{(2n)}(u) v^{2n}$$

near  $v = 0$ , where  $\gamma^{(j)} = d^j \gamma / dt^j$ . In particular, if we set  $H(u, v) := \sum_{n=0}^{\infty} \gamma^{(2n)}(u) v^n$ , then it gives a germ of real analytic function satisfying

$$H(u, -v^2) = \Phi(u, v), \quad H(u, v^2) = \Psi(u, v),$$

which prove that the images of  $\Phi$  and  $\Psi$  lie on the common real analytic surface. Since  $\gamma$  is non-degenerate, two vectors

$$H_u(u, 0) = \gamma'(u), \quad H_v(u, 0) = \gamma''(u)$$

are linearly independent, and  $H$  gives an immersion which contains  $\gamma$ .

Moreover, it can be easily checked that  $\Phi$  gives a space-like maximal surface (cf. Lemma 2.14) and  $\Psi$  gives a time-like minimal surface.

We now show the last assertion: Since the surface is real analytic, it is sufficient to show that given an arbitrary real analytic diffeomorphism  $f$  from  $(a, b)$  onto its image in  $\mathbf{R}$ ,

$$\Psi(u, v) = \frac{\gamma(u + iv) + \gamma(u - iv)}{2} \quad \text{and} \quad \tilde{\Psi}(u, v) = \frac{\tilde{\gamma}(u + iv) + \tilde{\gamma}(u - iv)}{2}$$

induce the same surface as their graphs, where  $\tilde{\gamma}(t) := \gamma(f(t))$ . We define  $A, B$  by

$$A = (f(u + v) + f(u - v))/2, \quad B = (f(u + v) - f(u - v))/2.$$

Thus it is sufficient to show that the map

$$(u, v) \mapsto (A, B)$$

is an immersion at  $(u, 0)$ . In fact, the Jacobian of the map is given by

$$J = \det \begin{pmatrix} f'(u) & 0 \\ 0 & f'(u) \end{pmatrix} \neq 0.$$

□

**Definition 2.3.** Let  $\Omega^2$  be a domain in  $\mathbf{R}^2$  and  $f : \Omega^2 \rightarrow \mathbf{R}$  a  $C^\infty$ -function satisfying (1.1). We set

$$B := 1 - f_x^2 - f_y^2.$$

A point  $p$  on  $\Omega^2$  is called a *non-degenerate point of type change*<sup>1</sup> with respect to  $f$  if

$$B(p) = 0, \quad \nabla B(p) \neq 0,$$

where  $\nabla B := (B_x, B_y)$  is the gradient vector of the function  $B$ .

Since  $\nabla B$  does not vanish at  $p$ , the function  $f$  actually changes type at the non-degenerate point  $p$ . The following assertion holds:

**Proposition 2.4.** *Let  $\gamma$  be a real analytic non-degenerate null curve, and let  $f_\gamma$  be the real analytic function induced by  $\gamma$  as in Proposition 2.2, which satisfies (1.1). Then the image of  $\gamma$  consists of non-degenerate points of type change with respect to  $f_\gamma$ .*

Note that the conclusion is stronger than that of Proposition 2.2.

*Proof.* Let  $\gamma$  be a non-degenerate null curve. Without loss of generality, we may take the time-component  $t$  as a parametrization of  $\gamma$ . Then we have the expression

$$\gamma(t) = (t, x(t), y(t)) \quad (a < t < b)$$

such that

$$(2.1) \quad x'(t)^2 + y'(t)^2 = 1.$$

Since  $\gamma$  is non-degenerate, it holds that

$$(2.2) \quad 0 \neq \gamma''(t) = (0, x''(t), y''(t)).$$

Differentiating the relation  $t = f(x(t), y(t))$ , we have that

$$(2.3) \quad x'(t)f_x(x(t), y(t)) + y'(t)f_y(x(t), y(t)) = 1.$$

On the other hand, the relation  $B = 0$  implies that

$$(2.4) \quad f_x(x(t), y(t))^2 + f_y(x(t), y(t))^2 = 1.$$

Then by (2.1), (2.3) and (2.4), it holds that

$$x'(t) = f_x, \quad y'(t) = f_y.$$

Thus we have that

$$(2.5) \quad \begin{aligned} (x'', y'') &= \frac{d}{dt} \left( f_x(x(t), y(t)), f_y(x(t), y(t)) \right) \\ &= (x'f_{xx} + y'f_{xy}, x'f_{xy} + y'f_{yy}) \\ &= (f_x f_{xx} + f_y f_{xy}, f_x f_{xy} + f_y f_{yy}) = -\frac{1}{2} \nabla B. \end{aligned}$$

By (2.2), we get the assertion. □

Conversely, we can prove the following.

---

<sup>1</sup> Klyachin [11] did not define this particular notion, but we include this definition here for purpose of clarification.

**Proposition 2.5** (Klyachin [11, Lemma 2]). *Let  $f : \Omega^2 \rightarrow \mathbf{R}$  be a  $C^\infty$ -function satisfying the zero mean curvature equation (1.1), and let  $p = (x_0, y_0) \in \Omega^2$  be a non-degenerate point of type change. Then there exists a non-degenerate  $C^\infty$ -regular null curve in  $\mathbf{R}_1^3$  with image passing through  $(f(x_0, y_0), x_0, y_0)$  and contained in the graph of  $f$ .*

*Proof.* By the implicit function theorem, there exists a unique  $C^\infty$ -regular curve  $\sigma(t) = (x(t), y(t))$  in the  $xy$ -plane with  $p = \sigma(0)$  so that  $B = 0$  along the curve. Since  $B = 0$  on  $\sigma$ , the velocity vector  $\sigma'$  is perpendicular to  $\nabla B$ . Since  $\nabla f$  is also perpendicular to  $\nabla B$ , we can conclude that  $\nabla f$  is proportional to  $\sigma'$ . In fact

$$\begin{aligned} -\frac{1}{2}\nabla f \cdot \nabla B &= (f_x, f_y) \begin{pmatrix} f_x f_{xx} + f_y f_{xy} \\ f_x f_{xy} + f_y f_{yy} \end{pmatrix} = f_x^2 f_{xx} + 2f_x f_y f_{xy} + f_y^2 f_{yy} \\ &= (1 - f_y^2) f_{xx} + 2f_x f_y f_{xy} + (1 - f_x^2) f_{yy} - (1 - f_x^2 - f_y^2)(f_{xx} + f_{yy}) \\ &= 0. \end{aligned}$$

Since  $f_x^2 + f_y^2 = 1$ , by taking an arclength parameter of  $\sigma$ , we may set

$$x' = f_x, \quad y' = f_y,$$

and then

$$B = 1 - f_x^2 - f_y^2 = 1 - (x')^2 - (y')^2 = 0$$

holds along  $\sigma$ , which implies that  $t \mapsto (t, x(t), y(t))$  is a null curve. Since

$$\frac{d}{dt} f(x(t), y(t)) = x' f_x + y' f_y = f_x^2 + f_y^2 = 1,$$

there exists a constant  $c$  such that  $f(x(t), y(t)) = t + c$ . By translating the graph vertically if necessary, we may assume that

$$f(x(t), y(t)) = t$$

holds for each  $t$ . Then we obtain the identity (2.5) in this situation, which implies that  $\nabla B(p) = (x''(0), y''(0)) \neq 0$ , namely,

$$(a, b) \ni t \mapsto (f(x(t), y(t)), x(t), y(t)) = (t, x(t), y(t)) \in \mathbf{R}_1^3$$

gives a non-degenerate null curve near  $t = 0$  lying in the graph of  $f$ .  $\square$

**Definition 2.6** ([2]). Let  $\Sigma^2$  be a Riemann surface. A  $C^\infty$ -map  $\varphi : \Sigma^2 \rightarrow \mathbf{R}_1^3$  is called a *generalized maximal surface* if there exists an open dense subset  $W$  of  $\Sigma^2$  such that the restriction  $\varphi|_W$  of  $\varphi$  to  $W$  gives a conformal (space-like) immersion of zero mean curvature. A *singular point* of  $\varphi$  is a point at which  $\varphi$  is not an immersion. A singular point  $p$  satisfying  $d\varphi(p) = 0$  is called a *branch point* of  $\varphi$ . Moreover,  $\varphi$  is called a *maxface* if  $\varphi$  does not have any branch point. (A maxface may have singular points in general).

*Remark 2.7.* The above definition of maxfaces is given in [2], which is simpler than the definition given in [13] and [5]. However, this new definition is equivalent to the previous one, as we now explain. Suppose that  $\varphi|_W$  is a conformal (space-like) immersion of zero mean curvature. Then  $\partial\varphi = \varphi_z dz$  is a  $\mathbf{C}^3$ -valued holomorphic 1-form on  $W$ , where  $z$  is a complex coordinate of  $\Sigma^2$ . Since  $\varphi$  is a  $C^\infty$ -map on  $\Sigma^2$ ,  $\partial\varphi$  can be holomorphically extended to  $\Sigma^2$ . Then the line integral  $\Phi(z) = \int_{z_0}^z \partial\varphi$  with respect to a base point  $z_0 \in \Sigma^2$  gives a holomorphic map defined on the universal cover of  $\Sigma^2$  whose real part coincides with  $\varphi(z) - \varphi(z_0)$ . The condition that  $\varphi$  does

not have any branch point implies that  $\Phi$  is an immersion. Moreover, since  $\varphi$  is conformal on  $W$ ,  $\Phi$  satisfies

$$-(d\Phi_0)^2 + (d\Phi_1)^2 + (d\Phi_2)^2 = 0 \quad (\Phi = (\Phi_0, \Phi_1, \Phi_2)),$$

namely  $\Phi$  is a null immersion. So  $\varphi$  satisfies the definition of maxface as in [13] and [5]. We call  $\Phi$  the *holomorphic lift* of the maxface  $\varphi$ .

Let  $\varphi : \Sigma^2 \rightarrow \mathbf{R}_1^3$  be a maxface with Weierstrass data  $(G, \eta)$  (see [13] for the definition of Weierstrass data). Using the data  $(G, \eta)$ , the maxface  $\varphi$  has the expression

$$(2.6) \quad \varphi = \operatorname{Re}(\Phi), \quad \Phi = \int_{z_0}^z (-2G, 1 + G^2, i(1 - G^2))\eta.$$

The imaginary part

$$(2.7) \quad \varphi^* := \operatorname{Im}(\Phi) : \tilde{\Sigma}^2 \longrightarrow \mathbf{R}_1^3$$

also gives a maxface called the *conjugate surface* of  $\varphi$ , which is defined on the universal cover  $\tilde{\Sigma}^2$  of  $\Sigma^2$ . The following fact is known:

**Fact 2.8** ([13, 5]). *A point  $p$  of  $\Sigma^2$  is a singular point of  $\varphi$  if and only if  $|G(p)| = 1$ .*

**Definition 2.9.** A singular point  $p$  of  $\varphi$  is called *non-degenerate* if  $dG$  does not vanish at  $p$ .

**Fact 2.10** ([13, 5]). *If a singular point  $p$  of  $\varphi$  is non-degenerate, then there exists a neighborhood  $U$  of  $p$  and a regular curve  $\gamma(t)$  in  $U$  so that  $\gamma(0) = p$  and the singular set of  $\varphi$  in  $U$  coincides with the image of the curve  $\gamma$ .*

This curve  $\gamma$  is called the *singular curve* at the non-degenerate singular point  $p$ .

**Definition 2.11.** A regular curve  $\gamma$  on  $\Sigma^2$  is called a *non-degenerate fold singularity* if it consists of non-degenerate singular points such that the real part of the meromorphic function  $dG/(G^2\eta)$  vanishes identically along the singular curve  $\gamma$ . Each point on the non-degenerate fold singularity is called a *fold singular point*.

A singular point of  $C^\infty$ -map  $\varphi : \Sigma^2 \rightarrow \mathbf{R}^3$  has a *fold singularity* at  $p$  if there exists a local coordinate  $(u, v)$  centered at  $p$  such that  $\varphi(u, v) = \varphi(u, -v)$ . Later, we show that a non-degenerate fold singularity is actually a fold singularity (cf. Lemma 2.15).

Suppose that  $p$  is a non-degenerate fold singular point of  $\varphi$ . The following duality between fold singularities and generalized cone-like singularities (cf. [4, Definition 2.1]) holds:

**Proposition 2.12** ([10]). *Let  $\varphi : \Sigma^2 \rightarrow \mathbf{R}_1^3$  be a maxface and  $\varphi^*$  the conjugate maxface. Then  $p$  is a non-degenerate fold singular point of  $\varphi$  if and only if it is a generalized cone-like singular point of  $\varphi^*$ .*

*Proof.* This assertion is immediate from comparison of the above definition of non-degenerate fold singularity and the definition of generalized cone-like singular points as in [4, Definition 2.1 and Lemma 2.3].  $\square$

We now show the following assertion, which characterizes the non-degenerate fold singularities on maxfaces.

**Theorem 2.13.** *Let  $\varphi : \Sigma^2 \rightarrow \mathbf{R}_1^3$  be a maxface which has non-degenerate fold singularities along a singular curve  $\gamma : (a, b) \rightarrow \Sigma^2$ . Then  $\hat{\gamma} := \varphi \circ \gamma$  is a non-degenerate null curve, and the image of the map*

$$(2.8) \quad \tilde{\varphi}(u, v) := \frac{\hat{\gamma}(u+v) + \hat{\gamma}(u-v)}{2}$$

*is real analytically connected to the image of  $\varphi$  along  $\gamma$  as a time-like minimal immersion. Conversely, any real analytic zero mean curvature immersion which changes type across a non-degenerate null curve is obtained as a real analytic extension of non-degenerate fold singularities of a maxface.*

This assertion follows immediately from Fact 2.2 and the following Lemmas 2.14 and 2.15.

**Lemma 2.14.** *Let  $\gamma : (a, b) \rightarrow \mathbf{R}_1^3$  be a real analytic non-degenerate null curve. Then*

$$\varphi(u+iv) := \frac{\gamma(u+iv) + \gamma(u-iv)}{2}$$

*gives a maxface with non-degenerate fold singularities on the real axis.*

*Proof.* We set  $z = u + iv$ . Then it holds that

$$\varphi_z = \frac{1}{2}(\varphi_u - i\varphi_v) = \frac{1}{2}\gamma'(u+iv),$$

where  $\gamma'(t) := d\gamma(t)/dt$ . Since  $\gamma$  is a regular real analytic curve, the map

$$(2.9) \quad \Phi(u+iv) := \gamma(u+iv)$$

gives a null holomorphic immersion if  $v$  is sufficiently small. Thus  $\varphi = \text{Re}(\Phi)$  gives a maxface.

Since  $\gamma$  is a null curve, it holds that

$$(2.10) \quad \gamma_0'(t)^2 = \gamma_1'(t)^2 + \gamma_2'(t)^2,$$

where we set  $\gamma = (\gamma_0, \gamma_1, \gamma_2)$ . Moreover, since  $\gamma$  is a regular curve, (2.10) implies

$$(2.11) \quad \gamma_0'(t) \neq 0 \quad (a < t < b).$$

It can be easily checked that the maxface  $\varphi$  has the Weierstrass data

$$(2.12) \quad \eta := \frac{1}{2}(d\Phi_1 - id\Phi_2) = \frac{\gamma_1'(z) - i\gamma_2'(z)}{2}dz,$$

$$(2.13) \quad g := -\frac{d\Phi_0}{2\eta} = -\frac{\gamma_0'(z)}{\gamma_1'(z) - i\gamma_2'(z)} = -\frac{\gamma_1'(z) + i\gamma_2'(z)}{\gamma_0'(z)},$$

where we set  $\Phi = (\Phi_0, \Phi_1, \Phi_2)$  and use the identity

$$(\gamma_1' - i\gamma_2')(\gamma_1' + i\gamma_2') = (\gamma_1')^2 + (\gamma_2')^2 = (\gamma_0')^2.$$

In particular, (2.13) implies that  $|G| = 1$  holds on the  $u$ -axis, which implies that the  $u$ -axis consists of singular points. By (2.13),  $dG$  vanishes if and only if

$$\Delta := (\gamma_1' + i\gamma_2')'\gamma_0' - (\gamma_1' + i\gamma_2')\gamma_0'' = (\gamma_0'\gamma_1'' - \gamma_1'\gamma_0'') + i(\gamma_0'\gamma_2'' - \gamma_2'\gamma_0'')$$

vanishes. In other words,  $\Delta = 0$  if and only if  $(\gamma_0'', \gamma_j'')$  is proportional to  $(\gamma_0', \gamma_j')$  for  $j = 1, 2$ , namely  $\gamma''$  is proportional to  $\gamma'$ . Since  $\gamma$  is non-degenerate, this is impossible. So the image of the curve  $\gamma$  consists of non-degenerate singular points (cf. Definition 2.9).

By (2.11), (2.12) and (2.13), we have that

$$\frac{dz}{G^2\eta} = \frac{2(\gamma'_1 - i\gamma'_2)}{(\gamma'_0)^2}.$$

Thus the  $u$ -axis consists of non-degenerate fold singular points if and only if  $\Delta_1 := (\gamma'_1 - i\gamma'_2)\Delta$  is a real valued function. Here

$$\operatorname{Re}(\Delta_1) = \gamma'_0(\gamma'_1\gamma''_1 + \gamma'_2\gamma''_2) - \gamma''_0((\gamma'_1)^2 + (\gamma'_2)^2) = \gamma'_0(\gamma'_0\gamma''_0) - \gamma''_0(\gamma'_0)^2 = 0,$$

where we used the identity (2.10) and its derivative. This implies that  $\gamma$  consists of non-degenerate fold singularities.  $\square$

Finally, we prove the converse assertion:

**Lemma 2.15.** *Let  $\varphi : \Sigma^2 \rightarrow \mathbf{R}_1^3$  be a maxface which has non-degenerate fold singularities along a singular curve  $\gamma : (a, b) \rightarrow \Sigma^2$ . Then, the space curve  $\hat{\gamma}(t) := \varphi \circ \gamma(t)$  is a non-degenerate real analytic null curve such that*

$$\hat{\varphi}(u, v) := \frac{1}{2} \left( \hat{\gamma}(u + iv) + \hat{\gamma}(u - iv) \right)$$

*coincides with the original maxface  $\varphi$ . In particular,  $\hat{\varphi}$  satisfies the identity  $\hat{\varphi}(u, v) = \hat{\varphi}(u, -v)$ .*

*Proof.* The singular set of  $\varphi$  can be characterized by the set  $|G| = 1$ , where  $(G, \eta)$  is the Weierstrass data as in (2.6). Let  $T$  be a Möbius transformation on  $S^2 = \mathbf{C} \cup \{\infty\}$  which maps the unit circle  $\{\zeta \in \mathbf{C} ; |\zeta| = 1\}$  to the real axis. Then  $T \circ G$  maps the image of the singular curve  $\gamma$  to the real axis. Since  $dG \neq 0$  (cf. Definition 2.11), we can choose  $T \circ G$  as a local complex coordinate. We denote it by

$$z = u + iv.$$

Then the image of  $\gamma$  coincides with the real axis  $\{v = 0\}$ . Let  $\Phi$  be the holomorphic lift of  $\varphi$ . Since the real axis consists of non-degenerate fold singularities, Proposition 2.12 implies that  $\operatorname{Im}(\Phi)$  is constant on the real axis. Since  $\Phi$  has an ambiguity of translations by pure imaginary vectors, we may assume without loss of generality that

$$(2.14) \quad \operatorname{Im}(\Phi) = 0 \quad \text{on the real axis.}$$

Thus, the curve  $\hat{\gamma}$  is expressed by (cf. (2.9))

$$(2.15) \quad \hat{\gamma}(u) = \varphi(u, 0) = \operatorname{Re}(\Phi(u, 0)) = \Phi(u, 0),$$

namely, two holomorphic functions  $\Phi(u + iv)$  and  $\hat{\gamma}(u + iv)$  take same values on the real axis. Hence  $\Phi(z) = \hat{\gamma}(u + iv)$  and thus

$$\varphi(z) = \frac{\hat{\gamma}(u + iv) + \hat{\gamma}(u - iv)}{2} = \hat{\varphi}(u, v).$$

So it is sufficient to show that  $\hat{\gamma}(t)$  is a non-degenerate null curve. Since  $|G| = 1$  on the real axis, there exists a real-valued function  $t = t(u)$  such that

$$(2.16) \quad G(u) = e^{it(u)} \quad (u \in \mathbf{R}).$$

Differentiating this along the real axis, we have  $G_u(u) = ie^{it}(dt/du)$ . Here,  $dt/du$  does not vanish because  $dG \neq 0$  on the real axis. Since  $\gamma$  consists of non-degenerate fold singularities,

$$i \frac{dG}{G^2\eta} = - \frac{e^{-it(u)}}{w(u)} \frac{dt}{du}$$

must be real valued (cf. Definition 2.11), where  $\eta = w(z) dz$ . Since  $\Phi$  is an immersion,  $G$  must have a pole at  $z = u$  if  $w(u) = 0$  (cf. (2.6)), but this contradicts the fact that  $|G| = 1$  along  $\gamma$ . Thus we have  $w(u) \neq 0$ . It then follows that

$$\xi(u) := e^{it(u)} w(u) = G(u) w(u)$$

is a non-vanishing real valued analytic function. Now, if we write  $\hat{\gamma} = (\hat{\gamma}_0, \hat{\gamma}_1, \hat{\gamma}_2)$ , (2.6) yields

$$\hat{\gamma}'_0(u) = \operatorname{Re} \left( -2G(u)w(u) \right) = -2\xi(u),$$

and

$$\hat{\gamma}'_1(u) = \operatorname{Re} \left( (1 + G(u)^2)w(u) \right) = 2\xi(u) \cos t(u),$$

$$\hat{\gamma}'_2(u) = \operatorname{Re} \left( i(1 - G(u)^2)w(u) \right) = 2\xi(u) \sin t(u).$$

This implies that  $\hat{\gamma}(u)$  is a regular real analytic null curve. Since  $dt/du \neq 0$ , the acceleration vector

$$\hat{\gamma}''(u) = (\log \xi(u))' \gamma'(u) + 2\xi(u) \begin{pmatrix} 0 \\ -\sin t(u) \\ \cos t(u) \end{pmatrix} \frac{dt}{du}$$

is not proportional to  $\hat{\gamma}'(u)$ . Thus,  $\hat{\gamma}(u)$  is non-degenerate.  $\square$

**Corollary 2.16.** *Let  $\varphi : \Sigma^2 \rightarrow \mathbf{R}_1^3$  be a maxface. Then a singular point  $p \in \Sigma^2$  lies on a non-degenerate fold singularity if and only if there exists a local complex coordinate  $z = u + iv$  with  $p = (0, 0)$  satisfying the following two properties:*

- (1)  $\varphi(u, v) = \varphi(u, -v)$ ,
- (2)  $\varphi_u(0, 0)$  and  $\varphi_{uu}(0, 0)$  are linearly independent.

*Proof.* Suppose  $p$  is a non-degenerate fold singularity of  $\varphi$ . Then (1) follows from Lemma 2.15, and (2) follows from the fact that the null curve parameterizing fold singularities is non-degenerate.

Conversely, suppose there is a coordinate system around a singular point  $p$  with  $p = (0, 0)$  which satisfies (1) and (2). Differentiating (1), we have that  $\varphi_v(u, 0) = 0$ . Let  $\Phi$  be a holomorphic lift of  $\varphi$ . Since  $\Phi$  is a null holomorphic map, the relation  $\varphi_v(u, 0) = 0$  implies that

$$0 = \langle \Phi_z(u, 0), \Phi_z(u, 0) \rangle = 4 \langle \varphi_z(u, 0), \varphi_z(u, 0) \rangle = \langle \varphi_u(u, 0), \varphi_u(u, 0) \rangle,$$

where  $\langle, \rangle$  is the canonical inner product of  $\mathbf{R}_1^3$ . This implies  $\gamma : u \mapsto \varphi(u, 0)$  is a null curve. By the condition (2), this null curve is non-degenerate. Then by Lemma 2.14,

$$\varphi_\gamma(u + iv) := \frac{\gamma(u + iv) + \gamma(u - iv)}{2}$$

is a maxface such that  $\gamma$  parametrizes a non-degenerate fold singularity of  $\varphi_\gamma$ . Moreover,  $\Phi_\gamma := \gamma(u + iv)$  gives a holomorphic lift of  $\varphi_\gamma$  (cf. (2.9)). Since

$$\Phi_z(u, 0) = 2\varphi_z(u, 0) = \gamma'(u) = (\Phi_\gamma)_z(u, 0),$$

the holomorphicity of  $\Phi$  and  $\Phi_\gamma$  yields that  $\Phi_z(z)$  coincides with  $(\Phi_\gamma)_z(z)$ . Thus  $\Phi$  coincides with  $\Phi_\gamma$  up to a constant. Then  $\varphi_\gamma$  coincides with  $\varphi$ , and we can conclude that  $\gamma$  parametrizes a non-degenerate fold singularity of  $\varphi$ .  $\square$

So far, we have looked at the singular curve of maxfaces. Now we turn our attention to the singular curve of zero mean curvature surfaces and prove the following assertion, which can be considered as the fundamental theorem of type change for zero mean curvature surfaces:

**Theorem 2.17.** *Let  $\gamma : (a, b) \rightarrow \mathbf{R}_1^3$  be a non-degenerate real analytic null curve. We set*

$$\hat{\varphi}_\gamma(u, v) := \begin{cases} \frac{\gamma(u + i\sqrt{v}) + \gamma(u - i\sqrt{v})}{2} & (v \geq 0), \\ \frac{\gamma(u + \sqrt{|v|}) + \gamma(u - \sqrt{|v|})}{2} & (v < 0), \end{cases}$$

for sufficiently small  $|v|$ . Then  $\hat{\varphi}_\gamma$  gives a real analytic zero mean curvature immersion such that the image of  $\gamma$  consists of non-degenerate points of type change with respect to  $\hat{\varphi}_\gamma$ .

Conversely, let  $f : \Omega^2 \rightarrow \mathbf{R}$  be a  $C^\infty$ -function satisfying the zero mean curvature equation (1.1), and let  $p = (x_0, y_0)$  be a non-degenerate point of type change with respect to  $f$ , where  $\Omega^2$  is a domain in the  $xy$ -plane. Then there exists a real analytic non-degenerate null curve  $\gamma$  in  $\mathbf{R}_1^3$  through  $(f(x_0, y_0), x_0, y_0)$  with  $\hat{\varphi}_\gamma$  coinciding with the graph of  $f$  in a small neighborhood of  $p$ .

Note that the conclusion for regularity of the converse statement is stronger than that of Proposition 2.5.

*Proof.* We have already proved the first assertion. (In fact, Proposition 2.4 implies that  $\gamma$  consists of non-degenerate points of type change with respect to  $\hat{\varphi}_\gamma$ .) Then it is sufficient to show the converse assertion, which is essentially due to Klyachin [11]. So, referring to [11], we give only a sketch of the proof: Let  $f : \Omega^2 \rightarrow \mathbf{R}$  be a  $C^\infty$ -function satisfying the zero mean curvature equation (1.1), and let  $p = (x_0, y_0) \in \Omega^2$  be a non-degenerate point of type change with respect to  $f$ . (One can prove the real analyticity of  $f$  at  $p$ , assuming only  $C^3$ -regularity of  $f$ , using the same argument as below.) By Proposition 2.5, there exists a  $C^\infty$ -regular curve  $\sigma(u)$  ( $|u| < \delta$ ) such that

$$\gamma(u) := (f \circ \sigma(u), \sigma(u))$$

is a non-degenerate null curve passing through  $(f(x_0, y_0), x_0, y_0)$  where  $\delta$  is a sufficiently small positive number. We set  $B := 1 - f_x^2 - f_y^2$ . Let  $\Omega^+$  be a simply connected domain such that  $B > 0$ , and suppose that  $\sigma$  lies on the boundary of  $\Omega^+$ . We set

$$t = t(x, y) := f(x, y), \quad s = s(x, y) := \int_{q_0}^q \frac{-f_y dx + f_x dy}{\sqrt{B}},$$

where  $q_0 \in \Omega^+$  is a base point and  $q := (x, y)$ . Since  $\alpha := (-f_y dx + f_x dy)/B$  is a closed 1-form, its (line) integral  $\int_{q_0}^q \alpha$  does not depend on our choice of path. Let  $\tau(v) = (a(v), b(v))$  ( $0 \leq v \leq \epsilon$ ) be a path starting from  $p$  and going into  $\Omega^+$  which is transversal to the curve  $\sigma$ . Since  $B(p) = 0$  and  $\nabla B(p) \neq 0$ , there exists a constant  $C > 0$  such that  $B \circ \tau(v) = Cv + \mathcal{O}(v^2)$ , where  $\mathcal{O}(v^2)$  denotes the higher

order terms. Then there exists a constant  $m$  such that

$$\left| \frac{-f_y \circ \tau(v) \frac{da}{dv}(v) + f_x \circ \tau(v) \frac{db}{dv}(v)}{\sqrt{B \circ \tau(v)}} \right| < \frac{m}{\sqrt{v}} \quad \text{for } 0 < v \leq \epsilon$$

hence

$$\lambda := \int_{\tau} |\alpha| < \int_0^{\epsilon} \frac{m}{\sqrt{v}} dv < \infty,$$

which is just the case (1) of [11, Lemma 6], and  $(t, s)$  gives an isothermal coordinate of  $\Omega^+$  with respect to the immersion

$$\varphi : (t, s) \mapsto (f(x(t, s), y(t, s)), x(t, s), y(t, s)) = (t, x(t, s), y(t, s)) \quad (s > 0).$$

Moreover, the function  $s(x, y)$  can be continuously extended to the image of the curve  $\sigma$ . Since  $\sigma$  is an integral curve of  $\nabla f$ , we may assume that  $\sigma$  parametrizes the level set  $s = 0$ , where we used the fact that  $s(x, y)$  is constant along each integral curve of  $\nabla f$ . In particular,  $\varphi$  satisfies  $\varphi_{tt} + \varphi_{ss} = 0$ . Then  $\varphi(t, s)$  can be extended to a harmonic  $\mathbf{R}^3$ -valued function for  $s < 0$  satisfying  $\varphi(t, s) = \varphi(t, -s)$  via the symmetry principle (see the proof of [11, Theorem 6]). In particular,  $f$  is a real analytic function whose graph coincides with the image of  $\varphi$  on  $\Omega^+$  near  $p$ . Moreover  $t \mapsto \varphi(t, 0)$  parametrizes the curve  $\gamma$  (cf. [11, Page 219]). By Corollary (2.16),  $\gamma$  can be considered as a non-degenerate fold singularity of the maxface  $(t, s) \mapsto \varphi(t, s)$ . (In fact, the condition (2) of Corollary (2.16) corresponds to the fact that  $\gamma$  is a non-degenerate curve near  $p$ .) Then Theorem 2.13 implies that  $\hat{\varphi}_\gamma$  coincides with the graph of  $f$  on a sufficiently small neighborhood of  $p$ .  $\square$

### 3. THE CONJUGATES OF HYPERBOLIC CATENOIDS AND SCHERK-TYPE SURFACES

The two entire graphs of  $n$  variables

$$f_1(x_1, \dots, x_n) := x_1 \tanh(x_2), \quad f_2(x_1, \dots, x_n) := (\log \cosh x_1) - (\log \cosh x_2),$$

given in Osamu Kobayashi [12], are zero mean curvature hypersurfaces in  $\mathbf{R}_1^{n+1}$  which change type from space-like to time-like. When  $n = 2$ , the image of  $f_1$  is congruent to  $\mathcal{C}_0$  and the image of  $f_2$  is congruent to  $\mathcal{S}_0$ . On the other hand, the space-like catenoid  $\mathcal{C}_+$  (resp. the space-like Scherk surface  $\mathcal{S}_+$ ) and the time-like catenoid  $\mathcal{C}_-$  (resp. the time-like Scherk surface  $\mathcal{S}_-$ ) are typical examples of zero mean curvature surfaces which contain singular light-like lines. Moreover, they are closely related to  $\mathcal{C}_0$  (resp.  $\mathcal{S}_0$ ) by taking their conjugate surfaces as follows:

**Proposition 3.1** ([8] and [3]). *The conjugate space-like maximal surface of the space-like hyperbolic catenoid  $\mathcal{C}_+$  and the conjugate time-like minimal surface of the time-like hyperbolic catenoid  $\mathcal{C}_-$  are both congruent to subsets of the entire graph  $\mathcal{C}_0$ .*

The space-like part of  $\mathcal{C}_0$  is connected and is called the *space-like hyperbolic helicoid*, and the time-like part of  $\mathcal{C}_0$  splits into two connected components, each of which is congruent to the *time-like hyperbolic helicoid* (see Figure 3, left).

*Proof.* This assertion has been pointed out in [8, Lemma 2.11 (3)] and the caption of Figure 1 in [3] without proof. We give a proof here as an application of the

results in the previous section. A subset of the space-like hyperbolic catenoid  $\mathcal{C}_+$  can be parametrized by

$$(3.1) \quad \begin{aligned} \varphi_1(u, v) &= (\cosh u \sin v, v, \sinh u \sin v) = -\operatorname{Re} i(\sinh z, z, \cosh z), \\ \psi_1(u, v) &= (-\cosh u \sin v, v, -\sinh u \sin v), \end{aligned}$$

where  $z = u + iv$ . In fact,  $\mathcal{C}_+$  is the union of the closure of the images of  $\varphi_1$  and  $\psi_1$ . The surface  $\varphi_1$  has generalized conical singularities at  $(u, n\pi)$  for any  $u \in \mathbf{R}$  and  $n \in \mathbf{Z}$ , as pointed out in [3]. Using this, one can easily compute that the conjugate of  $\varphi_1$  is congruent to the following surface

$$(3.2) \quad \begin{aligned} \varphi_1^*(u, v) &:= -\operatorname{Im} i(\sinh z, z, \cosh z) = -\operatorname{Re}(\sinh z, z, \cosh z) \\ &= -(\sinh u \cos v, u, \cosh u \cos v). \end{aligned}$$

By Proposition 2.12, the conjugate surface has non-degenerate fold singularities. Then by Theorem 2.13 one can get an analytic continuation of  $\varphi_1^*$  as a zero mean curvature surface in  $\mathbf{R}_1^3$  which changes type across the fold singularities. We can get an explicit description of such an extension of  $\varphi_1^*$  as follows: We set

$$(t, x, y) = \varphi_1^*(u, v) = -(\sinh u \cos v, u, \cosh u \cos v).$$

Then the surface has fold singularities at  $(u, n\pi)$  for any  $u \in \mathbf{R}$  and  $n \in \mathbf{Z}$ . Then it holds that

$$\frac{t}{y} = \tanh u = -\tanh x$$

and the image of  $-\varphi_1^*$  is contained in the surface  $\mathcal{C}_0$ .

On the other hand, a subset of the time-like hyperbolic catenoid  $\mathcal{C}_-$  has a parametrization

$$(3.3) \quad \varphi_2(u, v) := \frac{1}{2}(\sinh u + \sinh v, u + v, \cosh u - \cosh v) = \frac{\alpha(u) + \beta(v)}{2},$$

where

$$(3.4) \quad \alpha(u) := (\sinh u, u, \cosh u), \quad \beta(v) := (\sinh v, v, -\cosh v).$$

Also

$$(3.5) \quad \psi_2(u, v) := \frac{1}{2}(-\sinh u - \sinh v, u + v, -\cosh u + \cosh v)$$

gives a parametrization of  $\mathcal{C}_-$ . More precisely,  $\mathcal{C}_-$  is the union of the closure of the images of  $\varphi_2$  and  $\psi_2$ . We get the following parametrization of the conjugate surface  $\varphi_2^*$  of  $\varphi_2$

$$(3.6) \quad \varphi_2^*(u, v) := \frac{1}{2}(\alpha(u) - \beta(v)) = \frac{1}{2}(\sinh u - \sinh v, u - v, \cosh u + \cosh v),$$

where  $\alpha$  and  $\beta$  are as in (3.4). (See [7] for the definition of the conjugate surfaces of time-like minimal surfaces.) To find the implicit function representation of the image of  $\varphi_2^*$ , take a new coordinate system  $(\xi, \zeta)$  as

$$u = \xi + \zeta, \quad v = \xi - \zeta.$$

Then

$$\varphi_2^*(\xi, \zeta) = (\cosh \xi \sinh \zeta, \zeta, \cosh \xi \cosh \zeta),$$

which implies that the image is a subset of  $\mathcal{C}_0$ . The entire graph  $\mathcal{C}_0$  changes type across two disjoint real analytic null curves  $\{y = \pm \cosh x\}$ .  $\square$

Next, we prove a similar assertion for the Scherk surfaces, which is also briefly mentioned in the caption of Figure 1 in [3]:

**Theorem 3.2.** *The conjugate space-like maximal surface of the space-like Scherk surface  $\mathcal{S}_+$  and the conjugate time-like minimal surface of the time-like Scherk surface  $\mathcal{S}_-$  of the first kind are both congruent to subsets of the entire graph  $\mathcal{S}_0$ .*

*Proof.* Using the identities

$$\cos \arg z = \operatorname{Re}\left(\frac{z}{|z|}\right), \quad \sin \arg z = \operatorname{Im}\left(\frac{z}{|z|}\right),$$

one can prove that a subset of the space-like Scherk surface  $\mathcal{S}_+$  is parametrized by the complex variable  $z$  as

$$(3.7) \quad \begin{aligned} \varphi_1(z) &= -\operatorname{Re} i \left( \log \frac{1+z^2}{1-z^2}, \log \frac{1-z}{1+z}, \log \frac{1-iz}{1+iz} \right) + \frac{\pi}{2}(1, 1, 1) \\ &= \left( \arg \frac{1+z^2}{1-z^2}, \arg \frac{1-z}{1+z}, \arg \frac{1-iz}{1+iz} \right) + \frac{\pi}{2}(1, 1, 1). \end{aligned}$$

In fact,  $\varphi_1(z)$  is a multi-valued  $\mathbf{R}_1^3$ -valued function, but can be considered as a single-valued function on the universal covering of  $\mathbf{C} \cup \{\infty\} \setminus \{\pm 1, \pm i\}$ . We now set

$$\varphi_1(z) = (t(z), x(z), y(z))$$

and

$$\psi_1(z) := (\pi - t(z), x(z), \pi - y(z)).$$

Then  $\mathcal{S}_+$  is the union of the closure of the images of  $\varphi_1$  and  $\psi_1$ . The conjugate  $\psi_1^*$  of the space-like Scherk surface  $\psi_1$  is obtained by

$$(3.8) \quad \begin{aligned} \psi_1^*(z) &= \operatorname{Im} i \left( \log \frac{1+z^2}{1-z^2}, \log \frac{1-z}{1+z}, \log \frac{1-iz}{1+iz} \right) \\ &= \left( \log \left| \frac{1+z^2}{1-z^2} \right|, \log \left| \frac{1-z}{1+z} \right|, \log \left| \frac{1-iz}{1+iz} \right| \right). \end{aligned}$$

Since  $\psi_1$  admits only generalized cone-like singularities (cf. Proposition 2.12),  $\psi_1^*$  admits only fold singularities, and has a real analytical extension across the fold singularities to a time-like minimal surface in  $\mathbf{R}_1^3$  (cf. Theorem 2.13). More precisely, the image of the conjugate  $\psi_1^*$  is contained in the graph  $\mathcal{S}_0$ , as shown in the following: The singular sets of  $\psi_1$  and  $\psi_1^*$  are both parametrized as  $\{z = e^{iu}\}$ . Then the image of a connected component of singular curve by  $\psi_1^*$  as in (3.8) is parametrized as

$$(3.9) \quad \gamma(u) = \frac{1}{2} \left( 2 \log \cot u, \log \frac{1 - \cos u}{1 + \cos u}, \log \frac{1 + \sin u}{1 - \sin u} \right) \quad \left( 0 < u < \frac{\pi}{2} \right).$$

By the singular Björling formula (2.8) in Section 2, we have the following analytic extension of  $\gamma$

$$(3.10) \quad \hat{\psi}_2(u, v) := \frac{\gamma(u) + \gamma(v)}{2}.$$

Now, we check that the conjugate  $\psi_2 := \hat{\psi}_2^*$  of  $\hat{\psi}_2$  as in (3.10) coincides with the time-like Scherk surface  $\mathcal{S}_-$ . By (3.10), the conjugate  $\psi_2$  of  $\hat{\psi}_2$  is parametrized by

(see [7] for the definition of the conjugate surfaces of time-like minimal surfaces)

$$(3.11) \quad \begin{aligned} \psi_2(u, v) &= \frac{1}{2}(\gamma(u) - \gamma(v)) \\ &= \left( \frac{1}{2}(\log(\cot u) - \log(\cot v)), \frac{1}{4} \left( \log \frac{1 - \cos u}{1 + \cos u} - \log \frac{1 - \cos v}{1 + \cos v} \right), \right. \\ &\quad \left. \frac{1}{4} \left( \log \frac{1 + \sin u}{1 - \sin u} - \log \frac{1 + \sin v}{1 - \sin v} \right) \right). \end{aligned}$$

We set  $(t, x, y) = \psi_2(u, v)$ , and will show that  $(t, x, y)$  lies in  $\mathcal{S}_-$ : In fact, by (3.11), we have

$$e^{2t} = \frac{\cot u}{\cot v} = \frac{\cos u \sin v}{\sin u \cos v},$$

which implies

$$\cosh t = \frac{1}{2} \left( \sqrt{\frac{\cos u \sin v}{\sin u \cos v}} + \sqrt{\frac{\sin u \cos v}{\cos u \sin v}} \right) = \frac{\sin(u+v)}{\sqrt{\sin 2u \sin 2v}}.$$

Using

$$e^{4x} = \frac{1 - \cos u}{1 + \cos u} \times \frac{1 + \cos v}{1 - \cos v} = \left( \tan \frac{u}{2} \cot \frac{v}{2} \right)^2,$$

we have that

$$\cosh x = \frac{1}{\sqrt{\sin u \sin v}} \sin \frac{u+v}{2}.$$

Similarly,

$$\cosh y = \frac{1}{\sqrt{\cos u \cos v}} \cos \frac{u+v}{2}$$

holds. Hence the analytic extension of the conjugate of  $\mathcal{S}_+$  coincides with  $\mathcal{S}_0$ . As pointed out in [8], the space-like part of  $\mathcal{S}_0$  is connected, and the time-like part of  $\mathcal{S}_0$  consists of four connected components, each of which is congruent to the image of  $\psi_1$  (see Figure 3, right).  $\square$

#### 4. EXISTENCE OF ZERO MEAN CURVATURE SURFACES CHANGING TYPE ACROSS A LIGHT-LIKE LINE

We now discuss solutions of the zero mean curvature equation (1.1) which have the following form

$$(4.1) \quad f(x, y) = b_0(y) + \sum_{k=1}^{\infty} \frac{b_k(y)}{k} x^k,$$

where  $b_k(y)$  are  $C^\infty$ -functions. Since  $f$  contains a singular light-like line, we may assume without loss of generality that (cf. [3])

$$b_0(y) = y, \quad b_1(y) = 0.$$

As was pointed out in [3], there exists a real constant  $\mu$  called the *characteristic* of  $f$  such that  $b_2(y)$  satisfies the following equation

$$(4.2) \quad b_2'(y) + b_2(y)^2 + \mu = 0 \quad \left( ' = \frac{d}{dy} \right).$$

Now we derive the differential equations satisfied by  $b_k(y)$  for  $k \geq 3$ . If we set

$$Y := f_y - 1 = \sum_{k=2}^{\infty} \frac{b'_k}{k} x^k$$

and

$$\tilde{P} := 2(Yf_{xx} - f_x f_{xy}), \quad Q := Y^2 f_{xx} - 2f_x f_{xy} Y, \quad R := f_x^2 f_{yy},$$

then, by straightforward calculations, we see that

$$\begin{aligned} \tilde{P} &= -b_2 b'_2 x^2 - \frac{4}{3} b_2 b'_3 x^3 - \sum_{k=4}^{\infty} \left( P_k + \frac{2(k-1)}{k} b_2 b'_k + (3-k) b'_2 b_k \right) x^k, \\ Q &= - \sum_{k=4}^{\infty} Q_k x^k, \quad R = \sum_{k=4}^{\infty} R_k x^k. \end{aligned}$$

and that the zero mean curvature equation (1.1) reduces to

$$\sum_{k=2}^{\infty} \frac{b''_k}{k} x^k = f_{yy} = \tilde{P} + Q + R.$$

It is now immediate, by comparing the coefficients of  $x^k$  from both sides, to see that each  $b_k$  ( $k \geq 3$ ) satisfies the following ordinary differential equation

$$(4.3) \quad b''_k(y) + 2(k-1)b_2(y)b'_k(y) + k(3-k)b'_2(y)b_k(y) = -k(P_k + Q_k - R_k),$$

where  $P_3 = Q_3 = R_3 = 0$  and

$$\begin{aligned} P_k &:= \sum_{m=3}^{k-1} \frac{2(k-2m+3)}{k-m+2} b_m b'_{k-m+2}, \\ Q_k &:= \sum_{m=2}^{k-2} \sum_{n=2}^{k-m} \frac{3n-k+m-1}{mn} b'_m b'_n b_{k-m-n+2}, \\ R_k &:= \sum_{m=2}^{k-2} \sum_{n=2}^{k-m} \frac{b_m b_n b''_{k-m-n+2}}{k-m-n+2} \end{aligned}$$

for  $k \geq 4$ . Note that  $P_k$ ,  $Q_k$  and  $R_k$  are written in terms of  $b_j$  ( $j = 1, \dots, k-1$ ) and their derivatives.

Now, we consider the case that  $1 - f_x^2 - f_y^2$  changes sign across the light-like line  $t = y$ . This case occurs only when the characteristic  $\mu$  of  $f$  vanishes [3]. If we set

$$b_2(y) = 0 \quad (y \in \mathbf{R}),$$

then (4.2) holds for  $\mu = 0$ . So we assume

$$(4.4) \quad b_0(y) = y, \quad b_1(y) = 0, \quad b_2(y) = 0, \quad b_3(y) = 3cy,$$

where  $c$  is a non-zero constant. Then  $f(x, y)$  in (4.1) can be rewritten as

$$(4.5) \quad f(x, y) = y + cyx^3 + \sum_{k=4}^{\infty} \frac{b_k(y)}{k} x^k.$$

In this situation, we will find a solution satisfying

$$(4.6) \quad b_k(0) = b'_k(0) = 0 \quad (k \geq 4).$$




FIGURE 4. The graph of  $f(x, y)$  for  $c = 1/2$  and  $|x|, |y| < 0.8$  (The range of the graph is wider than the range used in our mathematical estimation. However, this figure still has a sufficiently small numerical error term in the Taylor expansion.)

Then (4.3) reduces to

$$(4.7) \quad b_k''(y) = -k(P_k + Q_k - R_k), \quad b_k(0) = b_k'(0) = 0, \quad (k = 4, 5, \dots),$$

$$(4.8) \quad P_k := \sum_{m=3}^{k-1} \frac{2(k-2m+3)}{k-m+2} b_m(y) b_{k-m+2}'(y) \quad (k \geq 4),$$

$$(4.9) \quad Q_k := \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{3n-k+m-1}{mn} b_m'(y) b_n'(y) b_{k-m-n+2}(y) \quad (k \geq 7),$$

$$(4.10) \quad R_k := \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{b_m(y) b_n(y) b_{k-m-n+2}''(y)}{k-m-n+2} \quad (k \geq 7),$$

and  $Q_k = R_k = 0$  for  $4 \leq k \leq 6$ , where the fact that  $b_2(y) = 0$  has been extensively used. For example,

$$\begin{aligned} b_0 &= y, & b_1 &= b_2 = 0, & b_3 &= 3cy, & b_4 &= -4c^2y^3, & b_5 &= 9c^3y^5, \\ b_6 &= -24c^2y^7, & b_7 &= 70c^5y^9 - 14c^3y^3, & \dots & \end{aligned}$$

In this section, we show the following assertion:

**Theorem 4.1.** *For each positive number  $c$ , the formal power series solution  $f(x, y)$  uniquely determined by (4.7), (4.8), (4.9) and (4.10) gives a real analytic zero mean*

curvature surface on a neighborhood of  $(x, y) = (0, 0)$ . In particular, there exists a non-trivial 1-parameter family of real analytic zero mean curvature surfaces each of which changes type across a light-like line (see Figure 4).

As a consequence, we get the following assertion:

**Corollary 4.2.** *There exists a family of zero-mean curvature hypersurfaces in Lorentz-Minkowski space  $\mathbf{R}_1^{n+1}$  which change type across an  $(n - 1)$ -dimensional light-like plane.*

*Proof.* Let  $f$  be as in the theorem. The graph of the function defined by

$$\mathbf{R}^n \ni (x_1, \dots, x_n) \mapsto f(x_1, x_2) \in \mathbf{R}$$

gives the desired hypersurface. In this case, the zero mean curvature equation

$$\left(1 - \sum_{j=1}^n f_{x_j}^2\right) \sum_{i=1}^n f_{x_i, x_i} + \sum_{i,j=1}^n f_{x_i} f_{x_j} = 0 \quad (f_{x_i} := \frac{\partial}{\partial x_i}, \quad f_{x_i, x_j} := \frac{\partial^2 f}{\partial x_i \partial x_j})$$

reduces to (1.1).  $\square$

To prove the theorem, it is sufficient to show that for arbitrary positive constants  $c > 0$  and  $\delta > 0$  there exist positive constants  $n_0$ ,  $\theta_0$ , and  $C$  such that

$$(4.11) \quad |b_k(y)| \leq \theta_0 C^k \quad (|y| \leq \delta)$$

holds for  $k \geq n_0$ . In fact, if (4.11) holds, then the series (4.5) converges uniformly over the rectangle  $[-C^{-1}, C^{-1}] \times [-\delta, \delta]$ .

The key assertion to prove (4.11) is the following

**Proposition 4.3.** *For each  $c > 0$  and  $\delta > 0$ , we set*

$$(4.12) \quad M := 3 \max \left\{ 144 c \tau |\delta|^{3/2}, \sqrt[4]{192 c^2 \tau} \right\},$$

where  $\tau$  is the positive constant given by (A.2) in the appendix, such that

$$(4.13) \quad t \int_t^{1-t} \frac{du}{u^2(1-u)^2} \leq \tau \quad \left(0 < t < \frac{1}{2}\right).$$

Then the function  $\{b_l(y)\}_{l \geq 3}$  formally determined by the recursive formula (4.7)–(4.10) satisfies the inequalities

$$(4.14) \quad |b_l''(y)| \leq c|y|^{l^*} M^{l-3},$$

$$(4.15) \quad |b_l'(y)| \leq \frac{3c|y|^{l^*+1}}{l^*+2} M^{l-3},$$

$$(4.16) \quad |b_l(y)| \leq \frac{3c|y|^{l^*+2}}{(l^*+2)^2} M^{l-3}$$

for any

$$(4.17) \quad y \in [-\delta, \delta],$$

where

$$(4.18) \quad l^* := \frac{1}{2}(l-1) - 2 \quad (l = 3, 4, \dots).$$

Once this proposition is proven, (4.11) follows immediately. In fact, if we set

$$\theta_0 = \frac{3}{c}(\delta M)^3, \quad C := \delta M$$

and  $n_0 \geq 7$ , then  $1 \leq l^* + 2 < l - 3$  and (4.11) follows from

$$\frac{3c|y|^{l^*+2}}{(l^*+2)^2} M^{l-3} \leq \theta_0 C^l.$$

## 5. PROOF OF PROPOSITION 4.3

We prove the proposition using induction on the number  $l \geq 3$ . If  $l = 3$ , then

$$\begin{aligned} |b_3''(y)| &= 0 \leq \frac{c}{|y|} = c|y|^{3^*} M^0, \\ |b_3'(y)| &= 3c = \frac{3c|y|^{3^*+1}}{3^*+2} M^0, \\ |b_3(y)| &= 3c|y| = \frac{3c|y|^{3^*+2}}{(3^*+2)^2} M^0 \end{aligned}$$

hold, using that  $b_3(y) = 3cy$ ,  $M^0 = 1$  and  $3^* = -1$ . So we prove the assertion for  $l \geq 4$ . Since (4.15), (4.16) follow from (4.14) by integration, it is sufficient to show that (4.14) holds for each  $l \geq 4$ . (In fact, the most delicate case is  $l = 4$ . In this case  $l^* = -1/2$  and we can use the fact that  $\int_0^{y_0} 1/\sqrt{y} dy$  for  $y_0 > 0$  converges.)

The inequality (4.14) follows if one shows that, for each  $k \geq 4$

$$(5.1) \quad |kP_k(y)|, |kQ_k(y)|, |kR_k(y)| \leq \frac{c}{3}|y|^{k^*} M^{k-3} \quad (|y| \leq \delta)$$

under the assumption that (4.14), (4.15) and (4.16) hold for all  $3 \leq l \leq k-1$ . In fact, if (5.1) holds, (4.14) for  $l = k$  follows immediately. Then by the initial condition (4.7) (cf. (4.6)), we have (4.15) and (4.16) for  $l = k$  by integration.

**The estimation of  $|kP_k|$  for  $k \geq 4$ .** By (4.8) and using the fact that (4.15), (4.16) hold for  $l \leq k-1$ , we have for each  $|y| < \delta$  that

$$\begin{aligned} |kP_k| &\leq \sum_{m=3}^{k-1} \frac{2k|k-2m+3|}{k-m+2} |b_m(y)| |b'_{k-m+2}(y)| \\ &\leq \sum_{m=3}^{k-1} \frac{2k|k-2m+3|}{k-m+2} \left( \frac{3cM^{m-3}|y|^{m^*+2}}{(m^*+2)^2} \right) \left( \frac{3cM^{k-m+2-3}|y|^{(k-m+2)^*+1}}{(k-m+2)^*+2} \right) \\ &= cM^{k-3}|y|^{k^*} \frac{144c|y|^{\frac{3}{2}}}{M} \sum_{m=3}^{k-1} \frac{k|k-2m+3|}{(m-1)^2(k-m+1)(k-m+2)} \\ &\leq cM^{k-3}|y|^{k^*} \frac{144c|\delta|^{\frac{3}{2}}}{M} \sum_{m=3}^{k-1} \frac{k|k-2m+3|}{(m-1)^2(k-m+1)(k-m+2)} \\ &\leq \frac{c}{3\tau} M^{k-3}|y|^{k^*} \sum_{m=3}^{k-1} \frac{k|k-2m+3|}{(m-1)^2(k-m+1)^2}. \end{aligned}$$

Here, we used (4.12). Since

$$\max_{m=3, \dots, k-1} |k-2m+3| = \max_{m=3, k-1} |k-2m+3| = \max\{|k-3|, |-k+5|\},$$

by setting  $q = m - 1$ , we have that

$$\begin{aligned} |kP_k| &\leq \frac{c}{3\tau} M^{k-3} |y|^{k^*} \sum_{m=3}^{k-1} \frac{k^2}{(m-1)^2 (k-m+1)^2} = \frac{c}{3\tau} M^{k-3} |y|^{k^*} \frac{1}{k} \sum_{q=2}^{k-2} \frac{k^3}{q^2 (k-q)^2} \\ &\leq \frac{c}{3\tau} M^{k-3} |y|^{k^*} \frac{1}{k} \int_{\frac{1}{k}}^{1-\frac{1}{k}} \frac{du}{u^2 (1-u)^2} \leq \frac{c}{3} M^{k-3} |y|^{k^*}, \end{aligned}$$

where we applied Lemma A.1 and (4.13) at the last step of the estimations. Hence, we get (5.1) for  $kP_k$ .

**The estimation of  $|kQ_k|$  for  $k \geq 7$ .** By (4.9) and the induction assumption, we have that

$$\begin{aligned} |kQ_k| &\leq \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{k|3n-k+m-1|}{mn} |b'_m(y)| |b'_n(y)| |b_{k-m-n+2}(y)| \\ &\leq \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{k|3n-k+m-1|}{mn} \left( \frac{3cM^{m-3}|y|^{m^*+1}}{m^*+2} \right) \times \\ &\quad \left( \frac{3cM^{n-3}|y|^{n^*+1}}{n^*+2} \right) \left( \frac{3cM^{k-m-n+2-3}|y|^{(k-m-n+2)^*+2}}{((k-m-n+2)^*+2)^2} \right) \\ &= cM^{k-3} |y|^{k^*} \frac{432c^2}{M^4} \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{k|3n-k+m-1|}{(m-1)^2 (n-1)^2 (k-m-n+2)^2}. \end{aligned}$$

Now we apply the inequality

$$\begin{aligned} \max_{\substack{3 \leq m \leq k-4 \\ 3 \leq n \leq k-m-1}} |3n-k+m-1| &= \max_{(m,n)=(3,3),(3,k-4),(k-4,3)} |3n-k+m-1| \\ &= \max\{|-k+11|, 4, |2k-10|\} \leq 2k, \end{aligned}$$

and also

$$\frac{432c^2}{M^4} \leq \frac{1}{36\tau},$$

which follows from (4.12). Setting  $p := m - 1$ ,  $q := n - 1$ , we have that

$$\begin{aligned} |kQ_k| &\leq \frac{c}{36\tau} M^{k-3} |y|^{k^*} \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{2k^2}{(m-1)^2 (n-1)^2 (k-m-n+2)^2} \\ &= \frac{c}{18\tau} M^{k-3} |y|^{k^*} \sum_{p=2}^{k-5} \sum_{q=2}^{k-p-2} \frac{k^2}{p^2 q^2 (k-p-q)^2}. \end{aligned}$$

Now applying Lemma A.2, we have that

$$|kQ_k| \leq \frac{c}{18\tau} M^{k-3} |y|^{k^*} \times 6\tau \leq \frac{c}{3} M^{k-3} |y|^{k^*},$$

which proves (5.1) for  $kQ_k$ .

**The estimation of  $|kR_k|$  for  $k \geq 7$ .** Like as in the case of  $|kQ_k|$ , we have that

$$\begin{aligned}
|kR_k| &\leq \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{k|b_m(y)| |b_n(y)| |b''_{k-m-n+2}(y)|}{k-m-n+2} \\
&\leq \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{k}{k-m-n+2} \left( \frac{3cM^{m-3}|y|^{m^*+2}}{(m^*+2)^2} \right) \times \\
&\quad \left( \frac{3cM^{n-3}|y|^{n^*+2}}{(n^*+2)^2} \right) \left( cM^{k-m-n+2-3}|y|^{(k-m-n+2)^*} \right) \\
&= 144c^3 M^{k-7} |y|^{k^*} \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{k}{(k-m-n+2)(m-1)^2(n-1)^2} \\
&= cM^{k-3} |y|^{k^*} \frac{144c^2}{M^4} \sum_{m=3}^{k-4} \sum_{n=3}^{k-m-1} \frac{k^2}{(k-m-n+2)^2(m-1)^2(n-1)^2}.
\end{aligned}$$

Now we set  $p = m - 1$ ,  $q = n - 1$ , and using the inequality

$$3^4 \times 144c^2\tau \leq 3^4 \times 192c^2\tau < M^4,$$

we have that

$$|kR_k| \leq \frac{c}{3^4\tau} M^{k-3} |y|^{k^*} \sum_{p=2}^{k-5} \sum_{q=2}^{k-p-2} \frac{k^2}{p^2 q^2 (k-p-q)^2}.$$

By applying Lemma A.2, we have that

$$|kR_k| \leq \frac{c}{3^4\tau} M^{k-3} |y|^{k^*} \times 6\tau < \frac{c}{3} M^{k-3} |y|^{k^*},$$

which proves (5.1) for  $kR_k$ . This completes the proof of Proposition 4.3.

#### APPENDIX A. INEQUALITIES USED IN SECTION 4

For  $a > 0$ , it holds that

$$(A.1) \quad \frac{1}{u^2(a-u)^2} = \frac{1}{a^3} \left( \frac{a}{u^2} + \frac{2}{u} + \frac{a}{(a-u)^2} + \frac{2}{a-u} \right).$$

Therefore,

$$(A.2) \quad \int_t^{a-t} \frac{du}{u^2(a-u)^2} = \frac{2}{a^3} \left( \frac{a(a-2t)}{t(a-t)} + 2 \log \frac{a-t}{t} \right) \quad (0 < t < \frac{a}{2}).$$

In particular, one can show that there exists a positive constant  $\tau$  such that

$$(A.3) \quad t \int_t^{1-t} \frac{du}{u^2(1-u)^2} \leq \tau \quad \left( 0 < t < \frac{1}{2} \right).$$

The following assertion is needed to prove (5.1) for  $kP_k(y)$ :

**Lemma A.1.** *Let  $p$  be a non-negative integer and  $k$  an integer satisfying  $k \geq p+4$ . Then the inequality*

$$\sum_{q=2}^{k-p-2} \frac{k^3}{q^2(k-p-q)^2} \leq \int_{\frac{1}{k}}^{a-\frac{1}{k}} \frac{du}{u^2(a-u)^2} \quad \left( a := 1 - \frac{p}{k} \right)$$

*holds.*

*Proof.* In fact, if we set  $a := 1 - (p/k)$ , then (A.1) yields that

$$\begin{aligned} \frac{k^3}{q^2(k-p-q)^2} &= \frac{1}{k} \frac{1}{\left(\frac{q}{k}\right)^2 \left(a - \frac{q}{k}\right)^2} \\ &= \frac{1}{a^3} \left[ \frac{1}{k} \left( \frac{a}{\left(\frac{q}{k}\right)^2} + \frac{2}{\left(\frac{q}{k}\right)} \right) + \frac{1}{k} \left( \frac{a}{\left(a - \frac{q}{k}\right)^2} + \frac{2}{a - \frac{q}{k}} \right) \right]. \end{aligned}$$

Since  $x \mapsto (a+2x)/x^2$  is a monotone decreasing function and the function  $x \mapsto (a+2(a-x))/(a-x)^2$  is monotone increasing on the interval  $(0, a/2)$ , we have that

$$\frac{k^3}{q^2(k-p-q)^2} \leq \frac{1}{a^3} \left[ \int_{\frac{q-1}{k}}^{\frac{q}{k}} \left( \frac{a}{u^2} + \frac{2}{u} \right) du + \int_{\frac{q}{k}}^{\frac{q+1}{k}} \left( \frac{a}{(a-u)^2} + \frac{2}{a-u} \right) du \right],$$

which yields that

$$\begin{aligned} \sum_{q=2}^{k-p-2} \frac{a^3 k^3}{q^2(k-p-q)^2} &\leq \int_{\frac{1}{k}}^{a-\frac{2}{k}} \left( \frac{a}{u^2} + \frac{2}{u} \right) du + \int_{\frac{2}{k}}^{a-\frac{1}{k}} \left( \frac{a}{(a-u)^2} + \frac{2}{a-u} \right) du \\ &\leq \int_{\frac{1}{k}}^{a-\frac{1}{k}} \left( \frac{a}{u^2} + \frac{2}{u} \right) du + \int_{\frac{1}{k}}^{a-\frac{1}{k}} \left( \frac{a}{(a-u)^2} + \frac{2}{a-u} \right) du \\ &\leq \int_{\frac{1}{k}}^{a-\frac{1}{k}} \left( \frac{a}{u^2} + \frac{2}{u} + \frac{a}{(a-u)^2} + \frac{2}{a-u} \right) du \\ &= \int_{\frac{1}{k}}^{a-\frac{1}{k}} \frac{du}{u^2(a-u)^2}. \end{aligned}$$

This proves the assertion.  $\square$

The following assertion is needed to prove (5.1) for  $kQ_k(y)$  and  $kR_k(y)$ :

**Lemma A.2.** *For any integer  $k \geq 7$ , the following inequalities hold:*

$$\sum_{p=2}^{k-5} \sum_{q=2}^{k-p-2} \frac{k^2}{p^2 q^2 (k-p-q)^2} \leq \frac{6}{k} \int_{\frac{1}{k}}^{1-\frac{1}{k}} \frac{du}{u^2(1-u)^2} \leq 6\tau,$$

where  $\tau$  is a constant satisfying (A.3).

*Proof.* We set  $a = a(p) := 1 - (p/k)$ . Applying Lemma A.1 and the identity (A.2), we have that

$$\begin{aligned} \sum_{p=2}^{k-5} \sum_{q=2}^{k-p-2} \frac{k^2}{p^2 q^2 (k-p-q)^2} &= \sum_{p=2}^{k-5} \left[ \frac{1}{kp^2} \sum_{q=2}^{k-p-2} \frac{k^3}{q^2(k-p-q)^2} \right] \\ &\leq \sum_{p=2}^{k-5} \left[ \frac{1}{kp^2} \int_{\frac{1}{k}}^{a-\frac{1}{k}} \frac{du}{u^2(a-u)^2} \right] = \sum_{p=2}^{k-5} \left[ \frac{1}{p^2} \frac{2}{a^2} \left( \frac{a-\frac{2}{k}}{a-\frac{1}{k}} + 2 \frac{\log(ka-1)}{ka} \right) \right] \\ &\leq \sum_{p=2}^{k-5} \left[ \frac{2}{p^2 a^2} \left( 1 + 2 \frac{\log ka}{ka} \right) \right] \leq \sum_{p=2}^{k-5} \frac{6}{p^2 a^2}, \end{aligned}$$

where we used the fact that  $\frac{\log ka}{ka} < 1$ . By applying Lemma A.1 and by using the property (A.3) of the constant  $\tau$ , it holds that

$$\begin{aligned} \sum_{p=2}^{k-5} \sum_{q=2}^{k-p-2} \frac{k^2}{p^2 q^2 (k-p-q)^2} &\leq 6 \sum_{p=2}^{k-5} \frac{1}{p^2 \left(1 - \frac{p}{k}\right)^2} = \frac{6}{k} \sum_{p=2}^{k-5} \frac{k^3}{p^2 (k-p)^2} \\ &\leq \frac{6}{k} \sum_{p=2}^{k-2} \frac{k^3}{p^2 (k-p)^2} \leq \frac{6}{k} \int_{\frac{1}{k}}^{1-\frac{1}{k}} \frac{du}{u^2(1-u)^2} < 6\tau, \end{aligned}$$

which proves the assertion.  $\square$

## REFERENCES

- [1] F. J. M. Estudillo and A. Romero, *Generalized maximal surfaces in Lorentz-Minkowski space  $L^3$* , Math. Proc. Camb. Phil. Soc., **111** (1992), 515–524.
- [2] S. Fujimori, Y. Kawakami, M. Kokubu, W. Rossman, M. Umehara, K. Yamada, *Hyperbolic Metrics on Riemann surfaces and Space-like CMC-1 surfaces in de Sitter 3-space*, M. Sánchez et al (eds), to appear in Recent Trends in Lorentzian Geometry, Springer Proceedings in Mathematics & Statistics 26.
- [3] S. Fujimori, Y.W. Kim, S.-E. Koh, W. Rossman, H. Shin, H. Takahashi, M. Umehara, K. Yamada and S.-D. Yang, *Zero mean curvature surfaces in  $L^3$  containing a light-like line*, to appear in C.R. Acad. Sci. Paris. Ser. I.
- [4] S. Fujimori, W. Rossman, M. Umehara, K. Yamada and S.-D. Yang, *New maximal surfaces in Minkowski 3-space with arbitrary genus and their cousins in de Sitter 3-space*, Results in Math. **56** (2009), 41–82.
- [5] S. Fujimori, K. Saji, M. Umehara and K. Yamada, *Singularities of maximal surfaces*, Math. Z., **259** (2008), 827–848.
- [6] C. Gu, *The extremal surfaces in the 3-dimensional Minkowski space*, Acta. Math. Sinica. **1** (1985), 173–180.
- [7] J. Inoguchi and M. Toda, *Timelike minimal surfaces via loop groups*, Acta. Appl. Math. **83** (2004), 313–335.
- [8] Y. W. Kim, S.-E. Koh, H. Shin and S.-D. Yang, *Spacelike maximal surfaces, timelike minimal surfaces, and Björling representation formulae*, Journal of Korean Math. Soc. **48** (2011), 1083–1100.
- [9] Y. W. Kim and S.-D. Yang, *A family of maximal surfaces in Lorentz-Minkowski three-space*, Proc. Amer. Math. Soc. **134** (2006), 3379–3390.
- [10] Y. W. Kim and S.-D. Yang, *Prescribing singularities of maximal surfaces via a singular Björling representation formula*, J. Geom. Phys., **57** (2007), 2167–2177.
- [11] V. A. Klyachin, *Zero mean curvature surfaces of mixed type in Minkowski space*, Izv. Math., **67** (2003), 209–224.
- [12] O. Kobayashi, *Maximal surfaces in the 3-dimensional Minkowski space  $L^3$* , Tokyo J. Math., **6** (1983), 297–309.
- [13] M. Umehara and K. Yamada, *Maximal surfaces with singularities in Minkowski space*, Hokkaido Math. J., **35** (2006), 13–40.

(Fujimori) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, OKAYAMA UNIVERSITY, OKAYAMA 700-8530, JAPAN

*E-mail address:* fujimori@math.okayama-u.ac.jp

(Kim) DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, SEOUL 136-701, KOREA

*E-mail address:* ywkim@korea.ac.kr

(Koh) DEPARTMENT OF MATHEMATICS, KONKUK UNIVERSITY, SEOUL 143-701, KOREA

*E-mail address:* sekoh@konkuk.ac.kr

(Rossman) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KOBE UNIVERSITY, KOBE 657-8501, JAPAN

*E-mail address:* wayne@math.kobe-u.ac.jp

(Shin) DEPARTMENT OF MATHEMATICS, CHUNG-ANG UNIVERSITY, SEOUL 156-756, KOREA  
*E-mail address:* `hshin@cau.ac.kr`

(Umehara) DEPARTMENT OF MATHEMATICAL AND COMPUTING SCIENCES, TOKYO INSTITUTE OF TECHNOLOGY, TOKYO 152-8552, JAPAN  
*E-mail address:* `umehara@is.titech.ac.jp`

(Yamada) DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, TOKYO 152-8551, JAPAN  
*E-mail address:* `kotaro@math.titech.ac.jp`

(Yang) DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, SEOUL 136-701, KOREA  
*E-mail address:* `sdyang@korea.ac.kr`